

Center of Infinitesimal Hecke Algebra of $\mathfrak{sl}_2$

Akaki Tikaradze

February 2, 2008

In this notes we are concerned with a special case of so called infinitesimal Hecke algebras introduced in [EGG] For simple Lie algebra $\mathfrak{sp}(2n)$ they construct certain algebra (called Infinitesimal Hecke algebra) which is endowed with a filtration so that corresponding associated graded algebra is isomorphic to the enveloping algebra of $\mathfrak{U}(\mathfrak{sp}(2n) \ltimes V)$, where V is an obvious representation (this is a version of PBW theorem). Following is the exact definition for the easiest case of $\mathfrak{sl}_2$.

Let $\mathbb{C}$ be the field of complex numbers, and let H_z be a semidirect product of $\mathfrak{U}\mathfrak{g}$ and TV (TV stands for tensor algebra), where by $\mathfrak{g}$ we will denote $\mathfrak{sl}_2$ and V is its two dimensional representation over $\mathbb{C}$ with basis x, y , mod out by the relation $[x, y] = z$, where z is an element of the center of $\mathfrak{U}\mathfrak{g}$. Thus H_z is an algebra depending on the parameter z , recall that $[e, y] = x$, and x is a maximal vector (killed by e) of weight 1 with respect to h (when talking about maximal vectors and such we always consider adjoint $\mathfrak{sl}_2$ action). Recall that one has an algebra filtration on H_z (x, y have degree 1, and $\mathfrak{U}\mathfrak{g}$ has degree 0), corresponding associated graded algebra is just $\mathfrak{U}(\mathfrak{g} \ltimes V)$ which also is equal to H_0 , this statement is a PBW theorem for this algebra (proved in [Kh]), and from now on we will denote this algebra just by H (suppressing the index 0). Representation theory of H_z was extensively studied in [Kh]. Here we will be concerned with the center and derivations of this algebra.

I would like to thank my advisor V. Ginzburg, and especially A. Khare for carefully going through all the computations and enormous help with use of Tex. Partially supported by NSF grant DMS-0401164.

Theorem 0.1. *Center of H_z is a polynomial algebra of one variable, generating central element is of the filtration degree 2.*

Proof. First, recall an anti-isomorphism of H_z which will be denoted by j , defined as: $j(x) = y, j(y) = x, j(h) = h, j(e) = -f, j(f) = -e$ ([Kh]). lets

note that this map fixes the following element $ey^2 + hxy - fx^2$. Indeed, applying this map to this element we get

$$-x^2f + xyh + y^2e = hxy + ey^2 - fx^2 - [e, y^2] - (-[f, x^2])$$

but $[e, y^2] = xy + yx = [f, x^2]$ so this element is indeed fixed, also since j is fixing h and center of $\mathfrak{U}\mathfrak{g}$, j is fixing the algebra generated by the above elements, h , and $\mathfrak{Z}(\mathfrak{U}\mathfrak{g})$ (center of $\mathfrak{U}L$). next step will be to exhibit an element from this algebra which will commute with e, x, h therefore will lie in the center of H_z . Let us denote the above element by $t = ey^2 + hxy - fx^2$, we have $[e, t] = e(xy + yx) - 2exy + hx^2 - hx^2 = -ez$, so we see that $[e, t - \frac{1}{2}hz] = 0$. We also have

$$\begin{aligned} [x, t] &= e(zx + xz) - x^2y + hxz + yx^2 \\ &= 2exz - e[z, y] + hzx - h[z, x] - (2zx - [z, x]) \\ &= 2exz - 2zx + [z, x] - e[z, y] + hzx - h[z, x], \end{aligned}$$

$$[x, \frac{1}{2}hz] = -\frac{1}{2}xz + \frac{1}{2}h[x, z] = -\frac{1}{2}zx + \frac{1}{2}[z, x] + \frac{1}{2}h[x, z].$$

Also,

$$[x, t - \frac{1}{2}hz] = 2exz - \frac{1}{2}3zx + \frac{1}{2}[z, x] - e[z, y] - \frac{1}{2}h[z, x] + hzx.$$

We denote the element $[x, t - \frac{1}{2}hz]$ by q_z , now we want to produce an element ω_z in the center of $\mathfrak{Z}(\mathfrak{U}L)$ so that $[x, \omega_z] = q_z$, then $t - \frac{1}{2}hz - \omega_z$ will be the desired central element. Let $\Delta = h^2 + 4ef - 2h$ be a multiple of the Casimir element, next we will compute commutators of powers of Δ with x and y , we have

$$[\Delta, x] = hx + xh + 4ey - 2x = (2h - 3)x + 4ey$$

$$[\Delta, y] = -hy - yh + 4xf + 2y = -(2hy + y) + 4(fx - y) + 2y = (-2h - 3)y + 4fx$$

Let us put $[\Delta^n, x] = (f_n(\Delta)h + g_n(\Delta))x + 2f_n(\Delta)ey$, (f_n, g_n are yet undetermined polynomials). It is always possible to write them in this way because $[\Delta^n, x]$ is a maximal vector of weight 1, so coefficient in front of y is maximal vector of weight 2, by classification of maximal vectors in $\mathfrak{U}\mathfrak{g}$ (explained below), this element is a product of e and an element of a center, similarly we treat coefficient in front of x . We have

$$\begin{aligned}
[\Delta^{n+1}, x] &= \Delta^n[\Delta, x] + [\Delta^n, x]\Delta \\
&= \Delta^n(2h - 3)x + \Delta^n 4ey + (f_n(\Delta)h + g_n(\Delta))x\Delta + 2f_n(\Delta)ey\Delta \\
&= (\Delta^n(2h - 3) + f_n(\Delta)\Delta h + g_n(\Delta)\Delta)x + (2f_n(\Delta)\Delta e + \Delta^n 4e)y \\
&\quad - (f_n(\Delta)h + g_n(\Delta))(2h - 3)x - 2f_n(\Delta)e4fx - \\
&\quad (f_n(\Delta)h + g_n(\Delta))4ey - 2f_n(\Delta)e(-2h - 3)y,
\end{aligned}$$

so grouping all elements containing y we get the following coefficient in front of y

$$\begin{aligned}
4\Delta^n e + 2f_n(\Delta)\Delta e - 4g_n(\Delta)e + 6f_n(\Delta)e + 4f_n(\Delta)(-2e) = \\
2(2\Delta^n + f_n(\Delta)(\Delta - 1) - 2g_n(\Delta))e
\end{aligned}$$

so $f_{n+1}(\Delta) = 2\Delta^n + f_n(\Delta)(\Delta - 1) - 2g_n(\Delta)$.

Grouping elements in front of x we get

$$\begin{aligned}
&\Delta^n(2h - 3) + f_n(\Delta)\Delta h + g_n(\Delta)\Delta - g_n(\Delta)2h + \\
&3g_n(\Delta) - f_n(\Delta)h + 4f_n(\Delta)h - 2f_n(\Delta)h^2 - 2f_n(\Delta)4ef,
\end{aligned}$$

sum of last 3 terms is $-2f_n(\Delta)\Delta$, so

$$\begin{aligned}
g_{n+1}(\Delta) &= -3\Delta^n + g_n(\Delta)\Delta + 3g_n(\Delta) - 2f_n(\Delta)\Delta = \\
&-3\Delta^n + (\Delta + 3)g_n(\Delta) - 2f_n(\Delta)\Delta.
\end{aligned}$$

We easily see that f_n is a polynomial of degree $n - 1$ with the top coefficient $2n$, in particular any equation $c = [z_1, x] + z_2 x$ has a unique solution in z_1, z_2 from $\mathfrak{Z}(\mathfrak{Ug})$, provided that c is of the form $2\psi ey + (h\psi + \psi_1)x$, where ψ, ψ_1 are also from $\mathfrak{Z}(\mathfrak{Ug})$ (because f_n form a basis of center, and by uniqueness in z_1 we mean uniqueness up to a constant), as an example of such c we can take arbitrary linear combinations of elements of the form $\Delta^i[\Delta^j, x]$. Remark that only elements from $\mathfrak{Z}(\mathfrak{Ug})$ which commute with x are constants. These observations will be needed later.

Recall that we want to write q_z as a commutator of x with a central element of $\mathfrak{Ug}$, let us rewrite q_z as follows:

$$\begin{aligned}
&z\left[\frac{1}{2}\Delta, x\right] - (e[z, y] + \frac{1}{2}h[z, x]) + \frac{1}{2}[z, x] = \\
&z\left[\frac{1}{2}\Delta, x\right] - [z, ey + \frac{1}{2}hx] + \frac{1}{2}[z, x] = z\left[\frac{1}{2}\Delta, x\right] - [z, [\frac{1}{4}\Delta, x]] - [z, \frac{3}{4}x] + \\
&\frac{1}{2}[z, x] = \frac{1}{4}(z[\Delta, x] + [\Delta, x]z - [z, x]).
\end{aligned}$$

Since $[\Delta z, x] = [\Delta, x]z + \Delta[z, x]$, its enough to show that there exists an element z_1 in the center of $\mathfrak{U}\mathfrak{g}$ so that $[z_1, x] = z[\Delta, x] - \Delta[z, x] = \Delta xz - zx\Delta$, let us take z_1, z_2 from center of $\mathfrak{U}\mathfrak{g}$ so that $z[\Delta, x] - \Delta[z, x] = [z_1, x] + z_2x$ (this is possible as remarked earlier). We want to show that $z_2=0$, applying $ad(f)$ to the last equality we get $z[\Delta, y] - \Delta[z, y] = [z_1, y] + z_2y$. Remark that $j(z[\Delta, x] - \Delta[z, x]) = -(z[\Delta, y] - \Delta[z, y])$, therefore

$$[z_1, y] + z_2y = -j([z_1, x] + z_2x) = [z_1, y] - yz_2 = [z_1 + z_2, y] - z_2y.$$

Applying $ad(e)$ to the last line of equalities and using uniqueness of z_2 implies that $z_2 = 0$, and we have constructed a central element $t_z = t - \frac{1}{2}hz - \omega_z$ of filtration degree 2. It remains to show that it generates the whole center. First claim is that any element which commutes with $\mathfrak{g}$ lies in the algebra generated by t_z and $\mathfrak{Z}(\mathfrak{U}\mathfrak{g})$.

Let a be such an element, write a as a polynomial in PBW basis (in x, y writing x on the right side of y with coefficients in $\mathfrak{U}\mathfrak{g}$), and let n be the smallest power of x appearing in this polynomial, so we can write $a = bx^n$ since a commutes with e and x commutes with e , implies that b has weight $-n$ and is maximal. However, since H_z is union of finite dimensional modules over $\mathfrak{g}$ (recall that action is always adjoint here) no maximal vectors can have negative weights, so $n = 0$. Let cy^m be a monomial of a with highest power of y appearing in monomials with no x in it, then $[e, a] = 0$ implies that $[e, c] = 0$ (because after taking commutator with e powers of x never decrease and power of y never increase), so c is a maximal vector of $\mathfrak{U}L$ of weight m . Now claim is that any maximal vector in $\mathfrak{U}\mathfrak{g}$ is generated by e, Δ . Indeed, let α be such a vector, using induction (assuming homogeneity) without loss of generality assume that it is not divisible by e , so if we write it in PBW basis (for $\mathfrak{U}\mathfrak{g}$) it will contain a monomial containing no e , so it has non positive weight so its weight is 0, therefore it lies in the center and we are done. So m is even $m = 2n$ and $c = \gamma e^{2n}$, γ being central, now let us consider an element $a - \gamma t_z^n$, call it ζ . Then ζ commutes with $\mathfrak{g}$ and has highest power of y without x less than m , now arguing inductively we will arrive at the central element of $\mathfrak{U}\mathfrak{g}$ so we are done. Finally, let a be a central element of H_z , in particular, it commutes with $\mathfrak{g}$ so it can be written as a polynomial of t_z with coefficients in $\mathfrak{Z}(\mathfrak{U}\mathfrak{g})$, let κt_z^n be the monomial of the top degree in a , since $[a, x] = 0$, passing to the associated graded ring we get that $[x, \kappa] = 0$, so κ is a constant, so we may disregard the top term, proceeding by induction we are done.

□

To summerise, $t_z = t - \frac{1}{2}hz - \omega_z$, where $t_z = ey^2 + hxy - fx^2$, and ω_z is a polynomial in Δ with the property $[x, \omega_z] = \frac{1}{4}([\Delta z - z, x] + z[\Delta, x] - \Delta[z, x])$. In what follows it will be convenient to use the following notation, denote by F, G linear endomorphisms of $C[\Delta]$ such that $F(\Delta^n) = f_n, G(\Delta^n) = g_n$. therefore F, G are surjective linear endomorphisms with kernel being constants. Thus in equation for ω_z after equating terms with ey we get

$$\frac{1}{4}(F(\Delta z - z) + zF(\Delta) - \Delta F(z)) = -F(\omega_z),$$

now lets rewrite recursive relations on f_n obtained before in terms of F, G , we get $F(\Delta z) = 2z + (\Delta - 1)F(z) - 2G(z)$, so $\omega_z = -\frac{1}{4}F^{-1}(F(\Delta z - z) + zF(\Delta) - \Delta F(z))$, thus

$$\begin{aligned}\omega_z &= -\frac{1}{4}(2z + F(z)(\Delta - 1) - 2G(z) - F(z) + 2z - \Delta F(z)) \\ &= -F^{-1}(z) + \frac{1}{2}z + \frac{1}{2}F^{-1}(G(z)),\end{aligned}$$

as a consequence of this formula, top coefficient of ω_z is $-\frac{1}{2n}$ times the top coefficient of z (assuming that the degree of z is $n-1$ as a polynomial of Δ), because according to computations of f_n, g_n , F and G decrease the degree of an input by 1, and top degree of f_n is $2n$.

If z is linear $z = a\Delta + b$, then explicit formula for a central element is the following:

$$ey^2 + hxy - fx^2 - \frac{1}{2}h(a\Delta + b) + \frac{1}{4}(a\Delta^2 + (2b - a)\Delta).$$

Theorem 0.2. *If z is non-zero then H_z has no outer derivations, and $H^1(H, H)$ is a rank 1 free module over the center of H .*

Proof. For the proof will need a description of maximal vectors of weight 1 in H . The claim is that this vector space as an algebra over the center of H_z is generated by $[x, \mathbb{C}[\Delta]]$ and by $\mathbb{C}[\Delta]x$. It suffices to show this when $z = 0$, let $r = \sum \alpha_{ij}x^i y^j$ be a vector of maximal weight, we will use induction on total degree with respect to x, y and biggest j so that α_{0j} is not zero. If such coefficients are all 0 then we can pull out x we will be left with a maximal vector of weight 0, so in that case nothing is to prove. We see that α_{0j} is maximal, so this term is equal to $\gamma e^n y^{2n-1}$, where γ is from $\mathbb{C}[\Delta]$, so $r - \gamma t^{n-1}(\frac{1}{4}[\Delta, x])$ is also a maximal vector whose biggest power of y without x is lower than j , so continuing this procedure all is left to show

is that $\alpha[\Delta, x]$ can be expressed appropriately, where α is a polynomial of Δ . Indeed, this element can be clearly written as $[\Delta, \gamma] + \omega x$, where ω, γ are polynomials of Δ (as explained earlier), so we are done.

We claim that $H^1(H, H)$ is rank 1 free module over the center, indeed assume that D is a derivation of H , we may assume that it vanishes on $\mathfrak{U}\mathfrak{g}$ (consequence of simplicity of the Lie algebra $\mathfrak{g}$), so D is a $\mathfrak{g}$ -map, therefore Dx is a maximal vector of rank 1, so by modifying with inner derivation, we may assume that $Dx = \sum t^i \alpha_i x$, where α_i is a polynomial of Δ . We claim that α_i is constant, indeed, assume the contrary, then

$$\begin{aligned} D(xy) &= \sum (t^i \alpha_i xy + xt^i \alpha_i y) = \sum t^i (2\alpha_i xy - [\alpha_i, x]y) \\ D(yx) &= \sum (t^i \alpha_i yx + yt^i \alpha_i x) = \sum t^i (2\alpha_i xy - [\alpha_i, y]x). \end{aligned}$$

Thus $\sum t^i [\alpha_i, x]y = \sum t^i [\alpha_i, y]x$, but this can not happen because after equating highest powers of t , left hand side contains terms with y^2 and right hand side does not, this implies that all α_i are constants, so D is given by $\mathfrak{g}$ -invariant Euler derivative on $\mathfrak{U}(\mathfrak{g} \ltimes V)$ (with respect to x, y) multiplied by a central element, therefore we are done. Now let's consider remaining cases (z non zero). We claim that every derivation is inner, suppose D is a derivation of H , again we may assume that it vanishes on $\mathfrak{U}\mathfrak{g}$. As before, we may assume that $Dx = \sum t_z^i \alpha_i x$, where α_i are central in $\mathfrak{U}\mathfrak{g}$ and α_0 is non zero, so $Dy = \sum t_z^i \alpha_i y$. Thus applying D to z we get

$$\begin{aligned} 0 = Dz &= [Dx, y] + [x, Dy] = 2 \sum t_z^i \alpha_i z + \sum t_z^i ([\alpha_i, y]x + [x, \alpha_i]y) \text{ so} \\ 2 \sum t_z^i \alpha_i z &= \sum t_z^i ([\alpha_i, x]y - [\alpha_i, y]x) \end{aligned}$$

For any α from $\mathbb{C}[\Delta]$, we have

$$\begin{aligned} &[\alpha, x]y - [\alpha, y]x \\ &= (2F(\alpha)ey + (F(\alpha)h + G(\alpha))x)y - (2F(\alpha)fx + (-F(\alpha)h + G(\alpha))y)x \\ &= 2F(\alpha)(ey^2 + hxy - fx^2 - \frac{1}{2}hz) + G(\alpha)z = 2F(\alpha)(t_z + \omega_z) + G(\alpha)z \end{aligned}$$

so our equality is as follows: $2 \sum t_z^i \alpha_i z = \sum t_z^i (2F(\alpha_i)(t + \omega_z) + G(\alpha_i)z)$, now let's equate terms with no t_z , we get $2\alpha_0 z = 2F(\alpha_0)B(z) + G(\alpha_0)z$. We have an equality in polynomials in Δ , so α_0 is not a constant otherwise right hand side would be 0, now comparing coefficients of top degrees in Δ gives a contradiction, because as we have observed F multiplies the top coefficient of an input by a positive number, however the top coefficient of ω_z is equal to a negative number times the top coefficient of z . End of the proof. $\square$

The University of Chicago, Department of Mathematics, Chicago, IL 60637, USA.

References

- [EGG] P. Etingof, W.L. Gan, and V. Ginzburg, *Continuous Hecke algebras*, Transform. Groups **10** (2005), no. 3-4, 423–447.
- [Kh] A. Khare, *Category $\mathcal{O}$ over a deformation of the symplectic oscillator algebra*, Journal of Pure and Applied Algebra, Volume **195**, No. 2 (2005), 131–166.